

TTC15 siRNA (h): sc-94264

BACKGROUND

The tetratricopeptide repeat (TPR) motif is a degenerate, 34 amino acid sequence found in many proteins and acts to mediate protein-protein interactions in various pathways. At the sequence level, there can be up to 16 tandem TPR repeats, each of which has a helix-turn-helix shape that stacks on other TPR repeats to achieve ligand binding specificity. TTC15 (TPR repeat protein 15), also known as TRAPPC12 (trafficking protein particle complex subunit 12) or CGI-87, is a 735 amino acid protein that contains four TPR repeats. Localizing to the endoplasmic reticulum (ER)-Golgi intermediate compartment, TTC15 may be involved in the early stage trafficking between the ER and Golgi apparatus, and is a component of the TRAPP tethering complex. The gene encoding TTC15 maps to human chromosome 2p25.3.

REFERENCES

1. Su, G., Roberts, T. and Cowell, J.K. 1999. TTC4, a novel human gene containing the tetratricopeptide repeat and mapping to the region of chromosome 1p31 that is frequently deleted in sporadic breast cancer. *Genomics* 55: 157-163.
2. Su, G., Casey, G. and Cowell, J.K. 2000. Genomic structure of the human tetratricopeptide repeat-containing gene, TTC4, from chromosome region 1p31 and mutation analysis in breast cancers. *Int. J. Mol. Med.* 5: 197-200.
3. Gauci, S., Helbig, A.O., Slijper, M., Krijgsvelde, J., Heck, A.J. and Mohammed, S. 2009. Lys-N and trypsin cover complementary parts of the phosphoproteome in a refined SCX-based approach. *Anal. Chem.* 81: 4493-4501.
4. Mayya, V., Lundgren, D.H., Hwang, S.I., Rezaul, K., Wu, L., Eng, J.K., Rodionov, V. and Han, D.K. 2009. Quantitative phosphoproteomic analysis of T cell receptor signaling reveals system-wide modulation of protein-protein interactions. *Sci. Signal.* 2: ra46.
5. Scrivens, P.J., Noueihed, B., Shahrzad, N., Hul, S., Brunet, S. and Sacher, M. 2011. C4orf41 and TTC-15 are mammalian TRAPP components with a role at an early stage in ER-to-Golgi trafficking. *Mol. Biol. Cell* 22: 2083-2093.
6. Wagner, S.A., Beli, P., Weinert, B.T., Nielsen, M.L., Cox, J., Mann, M. and Choudhary, C. 2011. A proteome-wide, quantitative survey of *in vivo* ubiquitylation sites reveals widespread regulatory roles. *Mol. Cell. Proteomics* 10: M111.013284.

CHROMOSOMAL LOCATION

Genetic locus: TRAPPC12 (human) mapping to 2p25.3.

PRODUCT

TTC15 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TTC15 shRNA Plasmid (h): sc-94264-SH and TTC15 shRNA (h) Lentiviral Particles: sc-94264-V as alternate gene silencing products.

For independent verification of TTC15 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-94264A, sc-94264B and sc-94264C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TTC15 siRNA (h) is recommended for the inhibition of TTC15 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TTC15 gene expression knockdown using RT-PCR Primer: TTC15 (h)-PR: sc-94264-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.