



# NOTO siRNA (h): sc-94333

## BACKGROUND

NOTO (notochord homeobox) is a 251 amino acid nuclear protein containing one homeobox DNA-binding domain. Noto is considered a transcription regulator that acts downstream of both HNF-3 $\beta$  and T during notochord development. Noto is required for node morphogenesis and for cilia formation in the posterior notochord (PNC). Essential for the expression of various components important for axonemal assembly and function, Noto plays an important role in regulating axial versus paraxial cell fate. Noto is encoded by a gene located on human chromosome 2p13.2. Chromosome 2 houses over 1,400 genes and comprises nearly 8% of the human genome. Harlequin ichthyosis, a rare and morbid skin deformity, is associated with mutations in the ABCA12 gene, while the lipid metabolic disorder sitosterolemia is associated with defects in the ABCG5 and ABCG8 genes. Additionally, an extremely rare recessive genetic disorder, Alström syndrome, is caused by mutations in the ALMS1 gene, which maps to chromosome 2.

## REFERENCES

1. Melby, A.E., Kimelman, D. and Kimmel, C.B. 1997. Spatial regulation of floating head expression in the developing notochord. *Dev. Dyn.* 209: 156-165.
2. Gritsman, K., Talbot, W.S. and Schier, A.F. 2000. Nodal signaling patterns the organizer. *Development* 127: 921-932.
3. Martinelli, C. and Spring, J. 2004. Expression pattern of the homeobox gene Not in the basal metazoan *Trichoplax adhaerens*. *Gene Expr. Patterns* 4: 443-447.
4. Abdelkhalik, H.B., Beckers, A., Schuster-Gossler, K., Pavlova, M.N., Burkhardt, H., Lickert, H., Rossant, J., Reinhardt, R., Schalkwyk, L.C., Müller, I., Herrmann, B.G., Ceolin, M., Rivera-Pomar, R. and Gossler, A. 2004. The mouse homeobox gene Not is required for caudal notochord development and affected by the truncate mutation. *Genes Dev.* 18: 1725-1736.
5. Beckers, A., Alten, L., Viebahn, C., Andre, P. and Gossler, A. 2007. The mouse homeobox gene Noto regulates node morphogenesis, notochordal ciliogenesis, and left right patterning. *Proc. Natl. Acad. Sci. USA* 104: 15765-15770.
6. Sand, F.W., Hörnblad, A., Johansson, J.K., Loren, C., Edsbacke, J., Stahlberg, A., Magenheimer, J., Ilavich, O., Mishani, E., Dor, Y., Ahlgren, U. and Semb, H. 2011. Growth-limiting role of endothelial cells in endoderm development. *Dev. Biol.* 352: 267-277.
7. Winzi, M.K., Hyttel, P., Dale, J.K. and Serup, P. 2011. Isolation and characterization of Node/Notochord-like cells from mouse embryonic stem cells. *Stem Cells Dev.* 20: 1817-1827.

## CHROMOSOMAL LOCATION

Genetic locus: Noto (human) mapping to 2p13.2.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## PRODUCT

NOTO siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Noto shRNA Plasmid (h): sc-94333-SH and Noto shRNA (h) Lentiviral Particles: sc-94333-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

NOTO siRNA (h) is recommended for the inhibition of Noto expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Noto gene expression knockdown using RT-PCR Primer: Noto (h)-PR: sc-94333-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.