

SLC4A1AP siRNA (h): sc-94489

BACKGROUND

Solute carrier family 4 anion exchanger member 1 adapter protein (SLC4A1AP), also designated Kanadaplin or human lung cancer oncogene 3 protein (HLC-3), is a 796 amino acid protein that is widely expressed in many tissues, including kidney, lung, liver, brain and skeletal and cardiac muscle. SLC4A1AP is a multidomain protein that localizes to the nucleus where it may play a role in signaling. SLC4A1AP was previously thought to act as an adaptor protein or chaperone involved in targeting kAE1 to the plasma membrane. However, recent studies suggest SLC4A1AP does not interact with kAE1. The gene encoding SLC4A1AP maps to chromosome 2, which consists of 237 million bases and makes up approximately 8% of the human genome. A number of genetic diseases are linked to genes on chromosome 2, including Harlequin ichthyosis, sitosterolemia and Alström syndrome.

REFERENCES

- Chen, J., et al. 1998. Kanadaplin is a protein that interacts with the kidney but not the erythroid form of band 3. *J. Biol. Chem.* 273: 1038-1043.
- Hübner, S., et al. 2002. Signal- and importin-dependent nuclear targeting of the kidney anion exchanger 1-binding protein kanadaplin. *Biochem. J.* 361: 287-296.
- Hübner, S., et al. 2003. Mitochondrial and nuclear localization of kanadaplin. *Eur. J. Cell Biol.* 82: 240-252.
- Kittanakom, S., et al. 2004. Human kanadaplin and kidney anion exchanger 1 (kAE1) do not interact in transfected HEK 293 cells. *Mol. Membr. Biol.* 21: 395-402.
- Hillier, L.W., et al. 2005. Generation and annotation of the DNA sequences of human chromosomes 2 and 4. *Nature* 434: 724-731.
- Thomas, A.C., et al. 2006. ABCA12 is the major harlequin ichthyosis gene. *J. Invest. Dermatol.* 126: 2408-2413.
- Marshall, J.D., et al. 2007. Alström syndrome. *Eur. J. Hum. Genet.* 15: 1193-1202.
- Marshall, J.D., et al. 2007. Spectrum of ALMS1 variants and evaluation of genotype-phenotype correlations in Alström syndrome. *Hum. Mutat.* 28: 1114-1123.

CHROMOSOMAL LOCATION

Genetic locus: SLC4A1AP (human) mapping to 2p23.3.

PRODUCT

SLC4A1AP siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SLC4A1AP shRNA Plasmid (h): sc-94489-SH and SLC4A1AP shRNA (h) Lentiviral Particles: sc-94489-V as alternate gene silencing products.

For independent verification of SLC4A1AP (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-94489A, sc-94489B and sc-94489C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SLC4A1AP siRNA (h) is recommended for the inhibition of SLC4A1AP expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

SLC4A1AP (49): sc-136069 is recommended as a control antibody for monitoring of SLC4A1AP gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SLC4A1AP gene expression knockdown using RT-PCR Primer: SLC4A1AP (h)-PR: sc-94489-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.