

SAP 14 siRNA (h): sc-94634

BACKGROUND

SAP 14 (spliceosome-associated protein, 14 kDa subunit), also known as P14, Ht006, CGI-110, HSPC175 or SF3B14a, is a 125 amino acid nuclear protein that is a component of the splicing factor 3b complex. Splicing factor 3b associates with both the U2 and U11/U12 small nuclear ribonucleoprotein complexes (U2 snRNP) of spliceosomes. Required for the splicing of pre-mRNA, SAP 14 enters the spliceosome and associates with the pre-mRNA branch site facilitating the interaction of snRNP with the branch sites of U2 and U12 of the 17S U2 and the 18S U11/U12 snRNP complex. SAP 14 contains a highly conserved RRM (RNA recognition motif) domain and interacts with SAP 155. SAP 14 is encoded by a gene located on human chromosome 2p23.3, which houses over 1,400 genes and comprises nearly 8% of the human genome.

REFERENCES

- MacMillan, A.M., et al. 1994. Dynamic association of proteins with the pre-mRNA branch region. *Genes Dev.* 8: 3008-3020.
- Query, C.C., et al. 1996. Three recognition events at the branch-site adenosine. *EMBO J.* 15: 1392-1402.
- Will, C.L., et al. 2001. A novel U2 and U11/U12 snRNP protein that associates with the pre-mRNA branch site. *EMBO J.* 20: 4536-4546.
- Golas, M.M., et al. 2005. Major conformational change in the complex SF3b upon integration into the spliceosomal U11/U12 di-snRNP as revealed by electron cryomicroscopy. *Mol. Cell* 17: 869-883.
- Cass, D.M., et al. 2006. The SF3b155 N-terminal domain is a scaffold important for splicing. *Biochemistry* 45: 10092-10101.
- Dybkov, O., et al. 2006. U2 snRNA-protein contacts in purified human 17S U2 snRNPs and in spliceosomal A and B complexes. *Mol. Cell. Biol.* 26: 2803-2816.
- Spadaccini, R., et al. 2006. Biochemical and NMR analyses of an SF3b155-p14-U2AF-RNA interaction network involved in branch point definition during pre-mRNA splicing. *RNA* 12: 410-425.
- Kühn-Hölsken, E., et al. 2007. Improved identification of enriched peptide RNA cross-links from ribonucleoprotein particles (RNPs) by mass spectrometry. *Nucleic Acids Res.* 35: e95.
- Kuwasaki, K., et al. 2008. Complex assembly mechanism and an RNA-binding mode of the human p14-SF3b155 spliceosomal protein complex identified by NMR solution structure and functional analyses. *Proteins* 71: 1617-1636.

CHROMOSOMAL LOCATION

Genetic locus: SF3B6 (human) mapping to 2p23.3.

PRODUCT

SAP 14 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SAP 14 shRNA Plasmid (h): sc-94634-SH and SAP 14 shRNA (h) Lentiviral Particles: sc-94634-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SAP 14 siRNA (h) is recommended for the inhibition of SAP 14 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SAP 14 gene expression knockdown using RT-PCR Primer: SAP 14 (h)-PR: sc-94634-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.