

TREML2 siRNA (h): sc-95293

BACKGROUND

TREML2 (triggering receptor expressed on myeloid cells-like 2), also known as C6orf76 or TLT2, is a 321 amino acid single-pass type I membrane protein that contains one Ig-like V-type domain. Expressed in cultured B cells, as well as in T cell leukemia and monocyte leukemia, TREML2 functions as a cell surface receptor that is thought to play a role in the innate and adaptive immune responses. The gene encoding TREML2 maps to a region on human chromosome 6 that is shared by proteins which contain Ig variable (IgV) domains. Chromosome 6 contains 170 million base pairs and comprises nearly 6% of the human genome. Deletion of a portion of the q arm of chromosome 6 is associated with early onset intestinal cancer, suggesting the presence of a cancer susceptibility locus. Additionally, porphyria cutanea tarda, Parkinson's disease, Stickler syndrome and a susceptibility to bipolar disorder are all associated with genes that map to chromosome 6.

REFERENCES

1. Allcock, R.J., Barrow, A.D., Forbes, S., Beck, S. and Trowsdale, J. 2003. The human TREM gene cluster at 6p21.1 encodes both activating and inhibitory single IgV domain receptors and includes NKp44. *Eur. J. Immunol.* 33: 567-577.
2. Online Mendelian Inheritance in Man, OMIM[™]. 2005. Johns Hopkins University, Baltimore, MD. MIM Number: 609715. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. King, R.G., Herrin, B.R. and Justement, L.B. 2006. Trem-like transcript 2 is expressed on cells of the myeloid/granuloid and B lymphoid lineage and is up-regulated in response to inflammation. *J. Immunol.* 176: 6012-6021.
4. Wiersinga, W.J., Veer, C.T., Wieland, C.W., Gibot, S., Hooibrink, B., Day, N.P., Peacock, S.J. and van der Poll, T. 2007. Expression profile and function of triggering receptor expressed on myeloid cells-1 during melioidosis. *J. Infect. Dis.* 196: 1707-1716.
5. Hashiguchi, M., Kobori, H., Ritprajak, P., Kamimura, Y., Kozono, H. and Azuma, M. 2008. Triggering receptor expressed on myeloid cell-like transcript 2 (TLT-2) is a counter-receptor for B7-H3 and enhances T cell responses. *Proc. Natl. Acad. Sci. USA* 105: 10495-10500.
6. Molloy, E.J. 2009. Triggering receptor expressed on myeloid cells (TREM) family and the application of its antagonists. *Recent. Pat. Antiinfect. Drug. Discov.* 4: 51-56.

CHROMOSOMAL LOCATION

Genetic locus: TREML2 (human) mapping to 6p21.1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

TREML2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TREML2 shRNA Plasmid (h): sc-95293-SH and TREML2 shRNA (h) Lentiviral Particles: sc-95293-V as alternate gene silencing products.

For independent verification of TREML2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-95293A, sc-95293B and sc-95293C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TREML2 siRNA (h) is recommended for the inhibition of TREML2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TREML2 gene expression knockdown using RT-PCR Primer: TREML2 (h)-PR: sc-95293-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.