

β-defensin 110 siRNA (h): sc-95324

BACKGROUND

β-defensins (also designated BDs, or hBDs in human) are small cationic peptides with broad-spectrum antimicrobial activity against a variety of enveloped viruses, fungi and bacteria. Produced in mucosal epithelia and neutrophils of several species, β-defensins are developmentally regulated. The family of β-defensin proteins share a common defensin-motif that is characterized by multiple cysteine residues and a highly conserved tertiary structure. Besides playing a significant role in host immune defense, many β-defensins also are involved in sperm maturation and capacitation. β-defensin 110 is a 67 amino acid secreted protein that has antibacterial activity. There are two isoforms of β-defensin 110 that exist as a result of alternative splicing events.

REFERENCES

1. Jia, H.P., Mills, J.N., Barahmand-Pour, F., Nishimura, D., Mallampali, R.K., Wang, G., Wiles, K., Tack, B.F., Bevins, C.L. and McCray, P.B. 1999. Molecular cloning and characterization of rat genes encoding homologues of human β-defensins. *Infect. Immun.* 67: 4827-4833.
2. Jia, H.P., Schutte, B.C., Schudy, A., Linzmeier, R., Guthmiller, J.M., Johnson, G.K., Tack, B.F., Mitros, J.P., Rosenthal, A., Ganz, T. and McCray, P.B. 2001. Discovery of new human β-defensins using a genomics-based approach. *Gene* 263: 211-218.
3. Kao, C.Y., Chen, Y., Zhao, Y.H. and Wu, R. 2003. ORFeome-based search of airway epithelial cell-specific novel human β-defensin genes. *Am. J. Respir. Cell Mol. Biol.* 29: 71-80.
4. Patil, A.A., Cai, Y., Sang, Y., Blecha, F. and Zhang, G. 2005. Cross-species analysis of the mammalian β-defensin gene family: presence of syntenic gene clusters and preferential expression in the male reproductive tract. *Physiol. Genomics* 23: 5-17.
5. Radhakrishnan, Y., Hamil, K.G., Yenugu, S., Young, S.L., French, F.S. and Hall, S.H. 2005. Identification, characterization, and evolution of a primate β-defensin gene cluster. *Genes Immun.* 6: 203-210.
6. Kouno, T., Fujitani, N., Mizuguchi, M., Osaki, T., Nishimura, S., Kawabata, S., Aizawa, T., Demura, M., Nitta, K. and Kawano, K. 2008. A novel β-defensin structure: a potential strategy of big defensin for overcoming resistance by Gram-positive bacteria. *Biochemistry* 47: 10611-10619.
7. Hosaka, Y., Koslowski, M., Nuding, S., Wang, G., Schlee, M., Schäfer, C., Saigenji, K., Stange, E.F. and Wehkamp, J. 2008. Antimicrobial host defense in the upper gastrointestinal tract. *Eur. J. Gastroenterol. Hepatol.* 20: 1151-1158.
8. Abedin, A., Mohammed, I., Hopkinson, A. and Dua, H.S. 2008. A novel antimicrobial peptide on the ocular surface shows decreased expression in inflammation and infection. *Invest. Ophthalmol. Vis. Sci.* 49: 28-33.
9. Diamond, G., Beckloff, N. and Ryan, L.K. 2008. Host defense peptides in the oral cavity and the lung: similarities and differences. *J. Dent. Res.* 87: 915-927.

CHROMOSOMAL LOCATION

Genetic locus: DEFB110 (human) mapping to 6p12.3.

PRODUCT

β-defensin 110 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see β-defensin 110 shRNA Plasmid (h): sc-95324-SH and β-defensin 110 shRNA (h) Lentiviral Particles: sc-95324-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

β-defensin 110 siRNA (h) is recommended for the inhibition of β-defensin 110 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.