



UROC28 siRNA (h): sc-95392

BACKGROUND

UROC28, also known as PBOV1 (prostate and breast cancer overexpressed 1) or UC28, is a 135 amino acid cytoplasmic and nuclear protein that is expressed in colon, prostate, small intestine, testis and spleen, with lower expression in thymus, ovary and peripheral blood leukocytes. UROC28 is up-regulated in prostate, breast and bladder cancer, and is thereby considered a useful target for diagnosis and treatment of these cancers. UROC28 is encoded by a gene located on human chromosome 6, which contains 170 million base pairs and comprises nearly 6% of the human genome. Deletion of a portion of the q arm of chromosome 6 is associated with early onset intestinal cancer, suggesting the presence of a cancer susceptibility locus. Additionally, Porphyria cutanea tarda, Parkinson's disease, Stickler syndrome and a susceptibility to bipolar disorder are all associated with genes that map to chromosome 6.

REFERENCES

1. An, G., Ng, A.Y., Meka, C.S., Luo, G., Bright, S.P., Cazares, L., Wright, G.L. and Veltri, R.W. 2000. Cloning and characterization of UROC28, a novel gene overexpressed in prostate, breast, and bladder cancers. *Cancer Res.* 60: 7014-7020.
2. Online Mendelian Inheritance in Man, OMIM™. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 605669. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Mungall, A.J., Palmer, S.A., Sims, S.K., Edwards, C.A., Ashurst, J.L., Wilming, L., Jones, M.C., Horton, R., Hunt, S.E., Scott, C.E., Gilbert, J.G., Clamp, M.E., Bethel, G., Milne, S., Ainscough, R., Almeida, J.P., et al. 2003. The DNA sequence and analysis of human chromosome 6. *Nature.* 425: 805-811.
4. Doak, S.H., Jenkins, S.A., Hurle, R.A., Varma, M., Hawizy, A., Kynaston, H.G. and Parry, J.M. 2007. Bone morphogenic factor gene dosage abnormalities in prostatic intraepithelial neoplasia and prostate cancer. *Cancer Genet. Cytogenet.* 176: 161-165.
5. Park, E., Kim, S., Kim, S.J., Park, Y., Lee, J.S., Yoo, J.C., Kim, C.S., Kim do, K., Lee, S.Y. and Chun, H.S. 2007. Modulation of parkin gene expression in noradrenergic neuronal cells. *Int. J. Dev. Neurosci.* 25: 491-497.
6. Bläker, H., Mechttersheimer, G., Sutter, C., Hertkorn, C., Kern, M.A., Rieker, R.J., Penzel, R., Schirmacher, P. and Kloor, M. 2008. Recurrent deletions at 6q in early age of onset non-HNPCC- and non-FAP-associated intestinal carcinomas. Evidence for a novel cancer susceptibility locus at 6q14-q22. *Genes Chromosomes Cancer* 47: 159-164.
7. Krukovskaia, L.L., Samusik, N.D., Shilov, E.S., Polev, D.E. and Kozlov, A.P. 2010. Tumor-specific expression of PBOV1, a new gene in evolution. *Vopr. Onkol.* 56: 327-332.

CHROMOSOMAL LOCATION

Genetic locus: PBOV1 (human) mapping to 6q23.3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

UROC28 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see UROC28 shRNA Plasmid (h): sc-95392-SH and UROC28 shRNA (h) Lentiviral Particles: sc-95392-V as alternate gene silencing products.

For independent verification of UROC28 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-95392A, sc-95392B and sc-95392C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

UROC28 siRNA (h) is recommended for the inhibition of UROC28 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor UROC28 gene expression knockdown using RT-PCR Primer: UROC28 (h)-PR: sc-95392-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.