



Ribosomal Protein S12 siRNA (h): sc-95463

BACKGROUND

Ribosomes, the organelles that catalyze protein synthesis, are composed of a small subunit (40S) and a large subunit (60S) that consist of over 80 distinct ribosomal proteins. Mammalian ribosomal proteins are encoded by multigene families that contain processed pseudogenes and one functional intron-containing gene within their coding regions. Ribosomal Protein S12 (RPS12) is a 132 amino acid component of the small ribosomal 40S subunit. The gene encoding Ribosomal Protein S12 is found on chromosome 6 which contains around 1,200 genes within 170 million base pairs of sequence. Deletion of a portion of the q arm of chromosome 6 is associated with early onset intestinal cancer suggesting the presence of a cancer susceptibility locus. Porphyria cutanea tarda is associated with chromosome 6 through the HFE gene which, when mutated, predisposes an individual to developing this porphyria.

REFERENCES

1. Wang, B., Malik, R., Nigg, E.A. and Körner, R. 2008. Evaluation of the low-specificity protease elastase for large-scale phosphoproteome analysis. *Anal. Chem.* 80: 9526-9533.
2. Daub, H., Olsen, J.V., Bairlein, M., Gnad, F., Oppermann, F.S., Körner, R., Greff, Z., Keri, G., Stemmann, O. and Mann, M. 2008. Kinase-selective enrichment enables quantitative phosphoproteomics of the kinome across the cell cycle. *Mol. Cell* 31: 438-448.
3. Dephoure, N., Zhou, C., Villen, J., Beausoleil, S.A., Bakalarski, C.E., Elledge, S.J. and Gygi, S.P. 2008. A quantitative atlas of mitotic phosphorylation. *Proc. Natl. Acad. Sci. USA* 105: 10762-10767.
4. Heibeck, T.H., Ding, S.J., Opresko, L.K., Zhao, R., Schepmoes, A.A., Yang, F., Tolmachev, A.V., Monroe, M.E., Camp, D.G., Smith, R.D., Wiley, H.S. and Qian, W.J. 2009. An extensive survey of tyrosine phosphorylation revealing new sites in human mammary epithelial cells. *J. Proteome Res.* 8: 3852-3861.
5. Mayya, V., Lundgren, D.H., Hwang, S.I., Rezaul, K., Wu, L., Eng, J.K., Rodionov, V. and Han, D.K. 2009. Quantitative phosphoproteomic analysis of T cell receptor signaling reveals system-wide modulation of protein-protein interactions. *Sci. Signal.* 2: ra46.
6. Doherty, L., Sheen, M.R., Vlachos, A., Choesmel, V., O'Donohue, M.F., Clinton, C., Schneider, H.E., Sieff, C.A., Newburger, P.E., Ball, S.E., Niewiadomska, E., Matysiak, M., Glader, B., Arcenci, R.J., et al. 2010. Ribosomal protein genes RPS10 and RPS26 are commonly mutated in Diamond-Blackfan anemia. *Am. J. Hum. Genet.* 86: 222-228.
7. Ren, J., Wang, Y., Liang, Y., Zhang, Y., Bao, S. and Xu, Z. 2010. Methylation of ribosomal protein S10 by protein-arginine methyltransferase 5 regulates ribosome biogenesis. *J. Biol. Chem.* 285: 12695-12705.
8. Burkard, T.R., Planyavsky, M., Kaupe, I., Breitwieser, F.P., Bürckstümmer, T., Bennett, K.L., Superti-Furga, G. and Colinge, J. 2011. Initial characterization of the human central proteome. *BMC Syst. Biol.* 5: 17.

CHROMOSOMAL LOCATION

Genetic locus: RPS12 (human) mapping to 6q23.2.

PRODUCT

Ribosomal Protein S12 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Ribosomal Protein S12 shRNA Plasmid (h): sc-95463-SH and Ribosomal Protein S12 shRNA (h) Lentiviral Particles: sc-95463-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Ribosomal Protein S12 siRNA (h) is recommended for the inhibition of Ribosomal Protein S12 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Ribosomal Protein S12 gene expression knockdown using RT-PCR Primer: Ribosomal Protein S12 (h)-PR: sc-95463-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.