

PGM 3 siRNA (h): sc-95517

BACKGROUND

Phosphoglucomutase (PGM), which belongs to the hexose-phosphate mutase family, plays an essential role in glycogen catabolism (glycogenolysis) as well as in the process of glycogen synthesis (glycogenesis). During glycogenolysis, PGM converts glucose-1-phosphate (Glc-1-P) to glucose-6-phosphate (Glc-6-P), thus promoting glycolysis and the pentose phosphate pathway. During glycogenesis, PGM functions in the opposite manner, converting glucose-6-phosphate into glucose-1-phosphate, to facilitate glycogen synthesis. PGM has five structural loci: PGM 1, PGM 2, PGM 3, PGM 4 and Aciculin. These five genetic forms of PGM differ in amino acid sequences but catalyze the same reactions, therefore indicating that they are isozymes. PGM 3 is a 542 amino acid protein expressed ubiquitously with the exception of lung tissue. Highest level of expression is found in heart, liver, pancreas and placenta tissue. All phosphoglucomutases act as monomers and bind one magnesium ion per subunit.

REFERENCES

- Lamm, L.U., et al. 1970. Linkage and association studies of two phosphoglucomutase loci (PGM1 and PGM3) to eighteen other markers. Analysis of the segregation at the marker loci. *Hum. Hered.* 20: 305-318.
- Takahashi, N., et al. 1982. A phylogeny for the principal alleles of the human phosphoglucomutase-1 locus. *Proc. Natl. Acad. Sci. USA* 79: 6636-6640.
- Whitehouse, D.B., et al. 1992. Phosphoglucomutase 1: complete human and rabbit mRNA sequences and direct mapping of this highly polymorphic marker on human chromosome 1. *Proc. Natl. Acad. Sci. USA* 89: 411-415.
- Pang, H., et al. 2002. Identification of human phosphoglucomutase 3 (PGM3) as N-acetylglucosamine-phosphate mutase (AGM1). *Ann. Hum. Genet.* 66: 139-144.
- Penha, L.L., et al. 2005. Cloning and characterization of the phosphoglucomutase of *Trypanosoma cruzi* and functional complementation of a *Saccharomyces cerevisiae* PGM null mutant. *Glycobiology* 15: 1359-1367.

CHROMOSOMAL LOCATION

Genetic locus: PGM3 (human) mapping to 6q14.1.

PRODUCT

PGM 3 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PGM 3 shRNA Plasmid (h): sc-95517-SH and PGM 3 shRNA (h) Lentiviral Particles: sc-95517-V as alternate gene silencing products.

For independent verification of PGM 3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-95517A, sc-95517B and sc-95517C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCL, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PGM 3 siRNA (h) is recommended for the inhibition of PGM 3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

PGM 3 (D-4): sc-390239 is recommended as a control antibody for monitoring of PGM 3 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PGM 3 gene expression knockdown using RT-PCR Primer: PGM 3 (h)-PR: sc-95517-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.