



EF-HC1 siRNA (h): sc-95518

BACKGROUND

EF-HC1 (EF-hand domain-containing protein 1), also known as Myoclonin-1, is a 640 amino acid protein containing an EF-hand domain and three DM10 domains. Interacting with the C-terminus of R-type Ca^{2+} CP $\alpha 1\text{E}$, EF-HC1 may enhance calcium influx and stimulate cell death. EF-HC1 is widely expressed in various tissues, excluding lymphocytes, and exists as three alternatively spliced isoforms. The gene encoding EF-HC1 maps to human chromosome 6p12.2. Mutations in the EF-HC1 gene have been linked to juvenile myoclonic epilepsy-1 (EJM1), a subtype of idiopathic generalized epilepsy with onset occurring during adolescence. EJM1 is characterized by afebrile seizures and myoclonic jerks, triggered by sleep deprivation, fatigue and alcohol consumption.

REFERENCES

1. Liu, A.W., et al. 1996. Juvenile myoclonic epilepsy in chromosome 6p12-p11: locus heterogeneity and recombinations. *Am. J. Med. Genet.* 63: 438-446.
2. Bai, D., et al. 2002. Juvenile myoclonic epilepsy: linkage to chromosome 6p12 in Mexico families. *Am. J. Med. Genet.* 113: 268-274.
3. Mungall, A.J., et al. 2003. The DNA sequence and analysis of human chromosome 6. *Nature* 425: 805-811.
4. Suzuki, T., et al. 2004. Mutations in EF-HC1 cause juvenile myoclonic epilepsy. *Nat. Genet.* 36: 842-849.
5. Gu, W., et al. 2005. A new EF-hand containing gene EF-HC2 on Xp11.4: tentative evidence for association with juvenile myoclonic epilepsy. *Epilepsy Res.* 66: 91-98.
6. Stogmann, E., et al. 2006. Idiopathic generalized epilepsy phenotypes associated with different EF-HC1 mutations. *Neurology* 67: 2029-2031.
7. Annesi, F., et al. 2007. Mutational analysis of EF-HC1 gene in Italian families with juvenile myoclonic epilepsy. *Epilepsia* 48: 1686-1690.
8. Bartocci, A., et al. 2007. Juvenile myoclonic epilepsy with generalised and focal electroencephalographic abnormalities: a case report with a molecular genetic study. *Neurol. Sci.* 28: 276-278.
9. Medina, M.T., et al. 2008. Novel mutations in Myoclonin-1/EF-HC1 in sporadic and familial juvenile myoclonic epilepsy. *Neurology* 70: 2137-2144.

CHROMOSOMAL LOCATION

Genetic locus: EFHC1 (human) mapping to 6p12.2.

PRODUCT

EF-HC1 siRNA (h) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see EF-HC1 shRNA Plasmid (h): sc-95518-SH and EF-HC1 shRNA (h) Lentiviral Particles: sc-95518-V as alternate gene silencing products.

For independent verification of EF-HC1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-95518A and sc-95518B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20°C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20°C , avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

EF-HC1 siRNA (h) is recommended for the inhibition of EF-HC1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor EF-HC1 gene expression knockdown using RT-PCR Primer: EF-HC1 (h)-PR: sc-95518-PR (20 μl). Annealing temperature for the primers should be $55-60^{\circ}\text{C}$ and the extension temperature should be $68-72^{\circ}\text{C}$.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.