

IPCEF1 siRNA (h): sc-95627

BACKGROUND

IPCEF1 (interaction protein for cytohesin exchange factors 1), also known as PIP3-E, is a 437 amino acid protein that localizes to both the cytoplasm and the cell membrane and contains one PH domain. Interacting with the guanine nucleotide exchange factors cytohesin-1, cytohesin-2, GRP1 and cytohesin-4, IPCEF1 enhances the guanine nucleotide exchange activity of Cytohesins with ARF6, thereby modifying ARF6 function and playing a role in Actin reorganization and membrane ruffling. Multiple isoforms of IPCEF1 exist due to alternative splicing events. The gene encoding IPCEF1 maps to human chromosome 6, which contains 170 million base pairs and comprises nearly 6% of the human genome. Deletion of a portion of the q arm of chromosome 6 is associated with early onset intestinal cancer, suggesting the presence of a cancer susceptibility locus. Additionally, porphyria cutanea tarda, Parkinson's disease, Stickler syndrome and a susceptibility to bipolar disorder are all associated with genes that map to chromosome 6.

REFERENCES

- Schürmann, A., Schmidt, M., Asmus, M., Bayer, S., Fliegert, F., Koling, S., Massmann, S., Schilf, C., Subauste, M.C., Voss, M., Jakobs, K.H. and Joost, H.G. 1999. The ADP-ribosylation factor (ARF)-related GTPase ARF-related protein binds to the ARF-specific guanine nucleotide exchange factor cytohesin and inhibits the ARF-dependent activation of phospholipase D. *J. Biol. Chem.* 274: 9744-9751.
- Kawada, N., Kristensen, D.B., Asahina, K., Nakatani, K., Minamiyama, Y., Seki, S. and Yoshizato, K. 2001. Characterization of a stellate cell activation-associated protein (STAP) with peroxidase activity found in rat hepatic stellate cells. *J. Biol. Chem.* 276: 25318-25323.
- Burmester, T., Ebner, B., Weich, B. and Hankeln, T. 2002. Cytochrome: a novel globin type ubiquitously expressed in vertebrate tissues. *Mol. Biol. Evol.* 19: 416-421.
- Krugmann, S., Anderson, K.E., Ridley, S.H., Risso, N., McGregor, A., Coadwell, J., Davidson, K., Eguinoa, A., Ellson, C.D., Lipp, P., Manfava, M., Ktistakis, N., Painter, G., Thuring, J.W., Cooper, M.A., Lim, Z.Y., et al. 2002. Identification of ARAP3, a novel PI3K effector regulating both ARF and Rho GTPases, by selective capture on phosphoinositide affinity matrices. *Mol. Cell* 9: 95-108.
- Venkateswarlu, K. 2003. Interaction protein for cytohesin exchange factors 1 (IPCEF1) binds cytohesin 2 and modifies its activity. *J. Biol. Chem.* 278: 43460-43469.
- Venkateswarlu, K. 2005. Analysis of the interaction between cytohesin 2 and IPCEF1. *Meth. Enzymol.* 404: 252-266.

CHROMOSOMAL LOCATION

Genetic locus: IPCEF1 (human) mapping to 6q25.2.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

IPCEF1 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see IPCEF1 shRNA Plasmid (h): sc-95627-SH and IPCEF1 shRNA (h) Lentiviral Particles: sc-95627-V as alternate gene silencing products.

For independent verification of IPCEF1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-95627A, sc-95627B and sc-95627C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

IPCEF1 siRNA (h) is recommended for the inhibition of IPCEF1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor IPCEF1 gene expression knockdown using RT-PCR Primer: IPCEF1 (h)-PR: sc-95627-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.