



RFX4 siRNA (h): sc-95773

BACKGROUND

EP and EP-like sites are regulatory enhancer elements found in the promoters of several viral and mammalian genes which, in humans, include the MIF-1 binding site (MIE) of the c-Myc gene, the X box of MHC class II promoters and a binding site in the PCNA (proliferating cell nuclear antigen) promoter. The EP-like sites present in the X box of MHC class II promoters are distinctly non-palindromic sequences that contain only a single EP-homologous half-site. The EP-like element is bound by a ubiquitous nuclear protein complex that consists of homo- and heterodimers involving the RFX1, RFX2, RFX3, RFX4 and RFX5 proteins. The RFX proteins represent an essential class II transcription factor family that share several conserved regions, including a centrally located DNA-binding domain (DBD) and a C-terminal D region that facilitates dimerization. RFX4 is a 735 amino acid nuclear protein that, via interactions with other RFX proteins, can bind DNA and is thought to activate the transcription of target genes. Four isoforms, each of which exhibit different tissue specificity, exist due to alternative splicing events.

REFERENCES

1. Reith, W., et al. 1994. RFX1, a transactivator of hepatitis B virus enhancer I, belongs to a novel family of homodimeric and heterodimeric DNA-binding proteins. *Mol. Cell. Biol.* 14: 1230-1244.
2. Emery, P., et al. 1996. RFX proteins, a novel family of DNA binding proteins conserved in the eukaryotic kingdom. *Nucleic Acids Res.* 24: 803-807.
3. Gajiwala, K.S., et al. 2000. Structure of the winged-helix protein hRFX1 reveals a new mode of DNA binding. *Nature* 403: 916-921.
4. Morotomi-Yano, K., et al. 2002. Human regulatory factor X 4 (RFX4) is a testis-specific dimeric DNA-binding protein that cooperates with other human RFX members. *J. Biol. Chem.* 277: 836-842.
5. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 603958. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Blackshear, P.J., et al. 2003. Graded phenotypic response to partial and complete deficiency of a brain-specific transcript variant of the winged helix transcription factor RFX4. *Development* 130: 4539-4552.
7. Araki, R., et al. 2004. Restricted expression and photic induction of a novel mouse regulatory factor X4 transcript in the suprachiasmatic nucleus. *J. Biol. Chem.* 279: 10237-10242.
8. Matsushita, H., et al. 2005. Identification of glioma-specific RFX4-E and -F isoforms and humoral immune response in patients. *Cancer Sci.* 96: 801-809.
9. Zhang, D., et al. 2008. G-protein pathway suppressor 2 (GPS2) interacts with the regulatory factor X4 variant 3 (RFX4_v3) and functions as a transcriptional co-activator. *J. Biol. Chem.* 283: 8580-8590.

CHROMOSOMAL LOCATION

Genetic locus: RFX4 (human) mapping to 12q23.3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

RFX4 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RFX4 shRNA Plasmid (h): sc-95773-SH and RFX4 shRNA (h) Lentiviral Particles: sc-95773-V as alternate gene silencing products.

For independent verification of RFX4 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-95773A, sc-95773B and sc-95773C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RFX4 siRNA (h) is recommended for the inhibition of RFX4 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RFX4 gene expression knockdown using RT-PCR Primer: RFX4 (h)-PR: sc-95773-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.