



PPM1H siRNA (h): sc-95791

BACKGROUND

PPM1H (protein phosphatase 1H), also known as URCC2 or ARHCL1, is a 514 amino acid protein that contains one PP2C-like domain and functions to catalyze the removal of phospho residues from target proteins. Belonging to the PP2C family, PPM1H can specifically dephosphorylate Cas, an event that may control the proliferation of cancer cells, suggesting an involvement for PPM1H in tumorigenesis. The gene encoding PPM1H maps to human chromosome 12, which encodes over 1,100 genes and comprises approximately 4.5% of the human genome. Chromosome 12 is associated with a variety of diseases and afflictions, including hypochondrogenesis, achondrogenesis, Kniest dysplasia, Noonan syndrome and Trisomy 12p, which causes facial developmental defects and seizure disorders.

REFERENCES

1. Allen, T.L., Brothman, A.R., Carey, J.C. and Chance, P.F. 1996. Cytogenetic and molecular analysis in trisomy 12p. *Am. J. Med. Genet.* 63: 250-256.
2. Labes, M., Roder, J. and Roach, A. 1998. A novel phosphatase regulating neurite extension on CNS inhibitors. *Mol. Cell. Neurosci.* 12: 29-47.
3. Zumkeller, W., Volleth, M., Muschke, P., Tönnies, H., Heller, A., Liehr, T., Wieacker, P. and Stumm, M. 2004. Genotype/phenotype analysis in a patient with pure and complete trisomy 12p. *Am. J. Med. Genet. A* 129A: 261-264.
4. Montpetit, A., Larose, J., Boily, G., Langlois, S., Trudel, N. and Sinnett, D. 2004. Mutational and expression analysis of the chromosome 12p candidate tumor suppressor genes in pre-B acute lymphoblastic leukemia. *Leukemia* 18: 1499-1504.
5. Segel, R., Peter, I., Demmer, L.A., Cowan, J.M., Hoffman, J.D. and Bianchi, D.W. 2006. The natural history of trisomy 12p. *Am. J. Med. Genet. A* 140: 695-703.
6. Stein, R. 2007. Genetics of Noonan syndrome—a new gene, and the search is still on. *Clin. Genet.* 72: 402-404.
7. Sugiura, T., Noguchi, Y., Sakurai, K. and Hattori, C. 2008. Protein phosphatase 1H, overexpressed in colon adenocarcinoma, is associated with CSE1L. *Cancer Biol. Ther.* 7: 285-292.
8. Shreeram, S. and Bulavin, D.V. 2008. PPM1H—new kid on the block. *Cancer Biol. Ther.* 7: 293-294.
9. Dephoure, N., Zhou, C., Villen, J., Beausoleil, S.A., Bakalarski, C.E., Elledge, S.J. and Gygi, S.P. 2008. A quantitative atlas of mitotic phosphorylation. *Proc. Natl. Acad. Sci. USA* 105: 10762-10767.

CHROMOSOMAL LOCATION

Genetic locus: PPM1H (human) mapping to 12q14.1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

PPM1H siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PPM1H shRNA Plasmid (h): sc-95791-SH and PPM1H shRNA (h) Lentiviral Particles: sc-95791-V as alternate gene silencing products.

For independent verification of PPM1H (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-95791A, sc-95791B and sc-95791C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCL, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PPM1H siRNA (h) is recommended for the inhibition of PPM1H expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PPM1H gene expression knockdown using RT-PCR Primer: PPM1H (h)-PR: sc-95791-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.