



Ribosomal Protein L41 siRNA (h): sc-95840

BACKGROUND

Ribosomes, the organelles that catalyze protein synthesis, are composed of a small subunit (40S) and a large subunit (60S) that consist of over 80 distinct ribosomal proteins. Mammalian ribosomal proteins are encoded by multigene families that contain processed pseudogenes and one functional intron-containing gene within their coding regions. Ribosomal Protein L41 (RPL41 or HG12) is a 25 amino acid cytoplasmic protein that is part of the large 60S ribosomal subunit complex. A member of the L41E family of ribosomal proteins, Ribosomal Protein L41 is thought to interact with the β subunit of protein kinase CKII and stimulates phosphorylation of DNA topoisomerase II α by CKII. Like most other ribosomal proteins, Ribosomal Protein L41 exists as multiple processed pseudogenes that are scattered throughout the genome.

REFERENCES

1. Bagni, C., Mariottini, P., Annesi, F. and Amaldi, F. 1993. Human ribosomal protein L4: cloning and sequencing of the cDNA and primary structure of the protein. *Biochim. Biophys. Acta* 1216: 475-478.
2. Kajikawa, S., Nakayama, H., Suzuki, M., Takashima, A., Murayama, O., Nishihara, M., Takahashi, M. and Doi, K. 1998. Increased expression of rat ribosomal protein L4 mRNA in 5-azacytidine-treated PC12 cells prior to apoptosis. *Biochem. Biophys. Res. Commun.* 252: 220-224.
3. Kenmochi, N., Kawaguchi, T., Rozen, S., Davis, E., Goodman, N., Hudson, T.J., Tanaka, T. and Page, D.C. 1998. A map of 75 human ribosomal protein genes. *Genome Res.* 8: 509-523.
4. Trifa, Y., Privat, I., Gagnon, J., Baeza, L. and Lerbs-Mache, S. 1998. The nuclear RPL4 gene encodes a chloroplast protein that co-purifies with the T7-like transcription complex as well as plastid ribosomes. *J. Biol. Chem.* 273: 3980-3985.
5. Yoshihama, M., Uechi, T., Asakawa, S., Kawasaki, K., Kato, S., Higa, S., Maeda, N., Minoshima, S., Tanaka, T., Shimizu, N. and Kenmochi, N. 2002. The human ribosomal protein genes: sequencing and comparative analysis of 73 genes. *Genome Res.* 12: 379-390.
6. Ueno, M., Nakayama, H., Kajikawa, S., Katayama, K., Suzuki, K. and Doi, K. 2002. Expression of ribosomal protein L4 (rpl4) during neurogenesis and 5-azacytidine (5AzC)-induced apoptotic process in the rat. *Histol. Histopathol.* 17: 789-798.
7. Yang, H., Henning, D. and Valdez, B.C. 2005. Functional interaction between RNA helicase II/Guac and ribosomal protein L4. *FEBS J.* 272: 3788-3802.

CHROMOSOMAL LOCATION

Genetic locus: RPL41 (human) mapping to 12q13.2.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Ribosomal Protein L41 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Ribosomal Protein L41 shRNA Plasmid (h): sc-95840-SH and Ribosomal Protein L41 shRNA (h) Lentiviral Particles: sc-95840-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Ribosomal Protein L41 siRNA (h) is recommended for the inhibition of Ribosomal Protein L41 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Ribosomal Protein L41 gene expression knockdown using RT-PCR Primer: Ribosomal Protein L41 (h)-PR: sc-95840-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.