CCDC41 siRNA (h): sc-95924



The Power to Question

BACKGROUND

The coiled-coil domain is a structural motif found in proteins that are involved in a diverse array of biological functions such as the regulation of gene expression, cell division, membrane fusion and drug extrusion and delivery. CCDC41 (coiled-coil domain containing 41) is a 693 amino acid protein that exists as two alternatively spliced isoforms. The gene encoding CCDC41 maps to hu-man chromosome 12, which encodes over 1,100 genes and comprises approximately 4.5% of the human genome. Chromosome 12 is associated with a variety of diseases and afflictions, including hypochondrogenesis, achondrogenesis, Kniest dysplasia, Noonan syndrome and trisomy 12p, which causes facial developmental defects and seizure disorders.

REFERENCES

- Allen, T.L., et al. 1996. Cytogenetic and molecular analysis in trisomy 12p. Am. J. Med. Genet. 63: 250-256.
- 2. Gilbert, F., et al. 2000. Disease genes and chromosomes: disease maps of the human genome. Chromosome 12. Genet. Test. 4: 319-333.
- Montgomery, K.T., et al. 2001. A high-resolution map of human chromosome 12. Nature 409: 945-946.
- Mason, J.M., et al. 2004. Coiled coil domains: stability, specificity, and biological implications. Chembiochem 5: 170-176.
- Ota, T., et al. 2004. Complete sequencing and characterization of 21,243 full-length human cDNAs. Nat. Genet. 36: 40-45.
- 6. Riaz, N., et al. 2005. Genomewide significant linkage to stuttering on chromosome 12. Am. J. Hum. Genet. 76: 647-651.
- Scherer, S.E., et al. 2006. The finished DNA sequence of human chromosome 12. Nature 440: 346-351.
- Liu, J., et al. 2006. A seven-helix coiled coil. Proc. Natl. Acad. Sci. USA 103: 15457-15462.

CHROMOSOMAL LOCATION

Genetic locus: CCDC41 (human) mapping to 12q22.

PRODUCT

CCDC41 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CCDC41 shRNA Plasmid (h): sc-95924-SH and CCDC41 shRNA (h) Lentiviral Particles: sc-95924-V as alternate gene silencing products.

For independent verification of CCDC41 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-95924A, sc-95924B and sc-95924C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNAse-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNAse-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CCDC41 siRNA (h) is recommended for the inhibition of CCDC41 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CCDC41 gene expression knockdown using RT-PCR Primer: CCDC41 (h)-PR: sc-95924-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com