



COX14 siRNA (h): sc-96045

BACKGROUND

The cytochrome c oxidase (COX) family of proteins function as the final electron donor in the respiratory chain to drive a proton gradient across the inner mitochondrial membrane, ultimately resulting in the production of water. The mammalian COX apoenzyme is a dimer, with each monomer consisting of 13 subunits, some of which are mitochondrial and some of which are nuclear. COX14 (cytochrome c oxidase assembly protein COX14), also known as C12orf62, is a 57 amino acid single-pass mitochondrial membrane protein that is implicated in the stability and assembly of the COX complex. Defects in the COX14 gene are associated with mitochondrial complex IV deficiency.

REFERENCES

1. Kennaway, N.G., et al. 1990. Isoforms of mammalian cytochrome c oxidase: correlation with human cytochrome c oxidase deficiency. *Pediatr. Res.* 28: 529-535.
2. Burke, P.V., et al. 1997. Effects of oxygen concentration on the expression of cytochrome c and cytochrome c oxidase genes in yeast. *J. Biol. Chem.* 272: 14705-14712.
3. Petruzzella, V., et al. 1998. Identification and characterization of human cDNAs specific to BCS1, PET112, SCO1, COX15, and COX11, five genes involved in the formation and function of the mitochondrial respiratory chain. *Genomics* 54: 494-504.
4. Shoubridge, E.A. 2001. Cytochrome c oxidase deficiency. *Am. J. Med. Genet.* 106: 46-52.
5. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 614478. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Zee, J.M. and Glerum, D.M. 2006. Defects in cytochrome oxidase assembly in humans: lessons from yeast. *Biochem. Cell Biol.* 84: 859-869.
7. Barrientos, A., et al. 2009. Suppression mechanisms of COX assembly defects in yeast and human: insights into the COX assembly process. *Biochim. Biophys. Acta* 1793: 97-107.
8. Weraarpachai, W., et al. 2012. Mutations in C12orf62, a factor that couples COX I synthesis with cytochrome c oxidase assembly, cause fatal neonatal lactic acidosis. *Am. J. Hum. Genet.* 90: 142-151.

CHROMOSOMAL LOCATION

Genetic locus: COX14 (human) mapping to 12q13.12.

PRODUCT

COX14 siRNA (h) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see COX14 shRNA Plasmid (h): sc-96045-SH and COX14 shRNA (h) Lentiviral Particles: sc-96045-V as alternate gene silencing products.

For independent verification of COX14 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-96045A and sc-96045B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

COX14 siRNA (h) is recommended for the inhibition of COX14 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor COX14 gene expression knockdown using RT-PCR Primer: COX14 (h)-PR: sc-96045-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.