



## RERG siRNA (h): sc-96069

### BACKGROUND

Members of the Ras superfamily of small GTP-binding proteins are critical mediators of diverse cell signaling pathways, including those leading to cell proliferation, cytoskeletal organization and secretion. RERG (Ras-like, estrogen-regulated, growth inhibitor) is a 199 amino acid protein that localizes to the cytoplasm and belongs to the Ras subfamily of small GTPases. Expressed in pancreas, liver, skin, lung, brain, kidney and heart tissue, RERG possesses intrinsic GTPase activity and is able to bind both GDP and GTP and, when overexpressed, may reduce overall tumorigenic potential. The gene encoding RERG maps to human chromosome 12, which encodes over 1,100 genes and comprises approximately 4.5% of the human genome. Chromosome 12 is associated with a variety of diseases and afflictions, including hypochondrogenesis, achondrogenesis, Kniest dysplasia, Noonan syndrome and trisomy 12p, which causes facial developmental defects and seizure disorders.

### REFERENCES

1. Bauer, C.K., Engeland, B., Wulfsen, I., Ludwig, J., Pongs, O. and Schwarz, J.R. 1998. RERG is a molecular correlate of the inward-rectifying K current in clonal rat pituitary cells. *Recept. Channels* 6: 19-29.
2. Finlin, B.S., Gau, C.L., Murphy, G.A., Shao, H., Kimel, T., Seitz, R.S., Chiu, Y.F., Botstein, D., Brown, P.O., Der, C.J., Tamanoi, F., Andres, D.A. and Perou, C.M. 2001. RERG is a novel Ras-related, estrogen-regulated and growth-inhibitory gene in breast cancer. *J. Biol. Chem.* 276: 42259-42267.
3. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 612664. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Wang, A.G., Fang, W., Han, Y.H., Cho, S.M., Choi, J.Y., Lee, K.H., Kim, W.H., Kim, J.M., Park, M.G., Yu, D.Y., Kim, N.S. and Lee, D.S. 2006. Expression of the RERG gene is gender-dependent in hepatocellular carcinoma and regulated by histone deacetyltransferases. *J. Korean Med. Sci.* 21: 891-896.
5. Key, M.D., Andres, D.A., Der, C.J. and Repasky, G.A. 2006. Characterization of RERG: an estrogen-regulated tumor suppressor gene. *Meth. Enzymol.* 407: 513-527.

### CHROMOSOMAL LOCATION

Genetic locus: RERG (human) mapping to 12p12.3.

### PRODUCT

RERG siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RERG shRNA Plasmid (h): sc-96069-SH and RERG shRNA (h) Lentiviral Particles: sc-96069-V as alternate gene silencing products.

For independent verification of RERG (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-96069A, sc-96069B and sc-96069C.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

### APPLICATIONS

RERG siRNA (h) is recommended for the inhibition of RERG expression in human cells.

### SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

### RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RERG gene expression knockdown using RT-PCR Primer: RERG (h)-PR: sc-96069-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.