

PLBD1 siRNA (h): sc-96121

BACKGROUND

PLBD1 (phospholipase B domain containing 1), also known as putative phospholipase B-like 1, LAMA-like protein 1 or lamina ancestor homolog 1, is a 553 amino acid protein belonging to the phospholipase B-like family. The gene that encodes PLBD1 maps to human chromosome 12, which encodes over 1,100 genes, contains 132 million bases and makes up approximately 4.5% of the human genome. A number of skeletal deformities are linked to chromosome 12, including hypochondrogenesis, achondrogenesis and Kniest dysplasia. Noonan syndrome, characterized by heart and facial developmental defects, is caused by a mutant form of the PTPN11 gene product, SH-PTP2. Chromosome 12 is also linked to a homeobox gene cluster, which encodes crucial transcription factors for morphogenesis, and the natural killer complex gene cluster that encodes C-type lectin proteins, which mediate the NK cell response to MHC I interaction.

REFERENCES

1. Yang, W. and Cole, W.G. 1998. Low basal transcripts of the COL2A1 collagen gene from lymphoblasts show alternative splicing of exon 12 in the Kniest form of spondyloepiphyseal dysplasia. *Hum. Mutat.* 1: S1-S2.
2. Trowsdale, J., Barten, R., Haude, A., Stewart, C.A., Beck, S. and Wilson, M.J. 2001. The genomic context of natural killer receptor extended gene families. *Immunol. Rev.* 181: 20-38.
3. Kelley, J., Walter, L. and Trowsdale, J. 2005. Comparative genomics of natural killer cell receptor gene clusters. *PLoS Genet.* 1: 129-139.
4. van der Burgt, I. 2007. Noonan syndrome. *Orphanet J. Rare Dis.* 2: 4.
5. Martinez, F.O. 2009. The transcriptome of human monocyte subsets begins to emerge. *J. Biol.* 8: 99.
6. Xu, S., Cai, L., Zhao, L., Douhan-Hakansson, L., Kristjánsson, G., Pauksen, K. and Venge, P. 2010. Tissue localization and the establishment of a sensitive immunoassay of the newly discovered human phospholipase B-precursor (PLB-P). *J. Immunol. Methods* 353: 71-77.
7. Yang, T., Meng, Y., Shi, H., Zhao, S., Wang, G. and Huang, S. 2010. Mutation analysis of PTPN11 gene in Noonan syndrome. *Zhonghua Yi Xue Yi Chuan Xue Za Zhi* 27: 554-558.

CHROMOSOMAL LOCATION

Genetic locus: PLBD1 (human) mapping to 12p13.1.

PRODUCT

PLBD1 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PLBD1 shRNA Plasmid (h): sc-96121-SH and PLBD1 shRNA (h) Lentiviral Particles: sc-96121-V as alternate gene silencing products.

For independent verification of PLBD1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-96121A, sc-96121B and sc-96121C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PLBD1 siRNA (h) is recommended for the inhibition of PLBD1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PLBD1 gene expression knockdown using RT-PCR Primer: PLBD1 (h)-PR: sc-96121-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.