



Rab 39 siRNA (h): sc-96292

BACKGROUND

The Ras-related superfamily of guanine nucleotide binding proteins includes the R-Ras, Rap, Ral/Rec and Rho/Rab subfamilies, all of which are thought to play an important role in either endocytosis or in biosynthetic protein transport. The process of transporting newly synthesized proteins from the endoplasmic reticulum (ER) to various stacks of the Golgi complex and to secretory vesicles involves the movement of carrier vesicles and requires Rab protein function. Rab proteins are also an integral part of endocytic pathways. Rab 39, also known as Rab 39A, is a 217 amino acid protein that belongs to the small GTPase superfamily. Rab 39 may be involved in vesicular trafficking.

REFERENCES

1. Chen, D., Guo, J., Miki, T., Tachibana, M. and Gahl, W.A. 1996. Molecular cloning of two novel Rab genes from human melanocytes. *Gene* 174: 129-134.
2. Somsel Rodman, J. and Wandinger-Ness, A. 2000. Rab GTPases coordinate endocytosis. *J. Cell Sci.* 113: 183-192.
3. Stenmark, H. and Olkkonen, V.M. 2001. The Rab GTPase family. *Genome Biol.* 2: REVIEWS3007.
4. Rodriguez-Gabin, A.G., Cammer, M., Almazan, G., Charron, M. and Larocca, J.N. 2001. Role of rRab 22B, an oligodendrocyte protein, in regulation of transport of vesicles from trans Golgi to endocytic compartments. *J. Neurosci. Res.* 66: 1149-1160.
5. Bao, X., Faris, A.E., Jang, E.K. and Haslam, R.J. 2002. Molecular cloning, bacterial expression and properties of Rab 31 and Rab 32. *Eur. J. Biochem.* 269: 259-271.
6. Chen, T., Han, Y., Yang, M., Zhang, W., Li, N., Wan, T., Guo, J. and Cao, X. 2003. Rab39, a novel Golgi-associated Rab GTPase from human dendritic cells involved in cellular endocytosis. *Biochem. Biophys. Res. Commun.* 303: 1114-1120.
7. Ng, E.L., Wang, Y. and Tang, B.L. 2007. Rab 22B's role in *trans*-Golgi network membrane dynamics. *Biochem. Biophys. Res. Commun.* 361: 751-757.
8. Lodhi, I.J., Chiang, S.H., Chang, L., Vollenweider, D., Watson, R.T., Inoue, M., Pessin, J.E. and Saltiel, A.R. 2007. Gapex-5, a Rab 31 guanine nucleotide exchange factor that regulates Glut4 trafficking in adipocytes. *Cell Metab.* 5: 59-72.
9. Kotsch, M., Sieuwerts, A.M., Grosser, M., Meye, A., Fuessel, S., Meijer-van Gelder, M.E., Smid, M., Schmitt, M., Baretton, G., Luther, T., Magdolen, V. and Foekens, J.A. 2008. Urokinase receptor splice variant uPAR-del4/5-associated gene expression in breast cancer: identification of Rab 31 as an independent prognostic factor. *Breast Cancer Res. Treat.* 111: 229-240.

CHROMOSOMAL LOCATION

Genetic locus: RAB39A (human) mapping to 11q22.3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Rab 39 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Rab 39 shRNA Plasmid (h): sc-96292-SH and Rab 39 shRNA (h) Lentiviral Particles: sc-96292-V as alternate gene silencing products.

For independent verification of Rab 39 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-96292A, sc-96292B and sc-96292C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Rab 39 siRNA (h) is recommended for the inhibition of Rab 39 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Rab 39 gene expression knockdown using RT-PCR Primer: Rab 39 (h)-PR: sc-96292-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.