

PSMD13 siRNA (h): sc-96416

BACKGROUND

In eukaryotic cells, selective breakdown of cellular proteins is ensured by their ubiquitination and subsequent degradation by the 26S Proteasome. The 26S Proteasome is a protease complex that selectively breaks down proteins that have been modified by polyubiquitin chains. It is made up of two multi-subunit complexes: the 20S Proteasome chamber, which serves as the proteolytic core of the complex and two 19S regulatory particles which recognize and unfold ubiquitinated proteins. PSMD13 (proteasome (prosome, macropain) 26S subunit, non-ATPase 13), also known as S11, Rpn9, p40.5, or HSPC027, is a 376 amino acid protein that belongs to the proteasome subunit S11 family. PSMD13 acts as a regulatory subunit of the 26S proteasome which is involved in the ATP-dependent degradation of ubiquitinated proteins.

REFERENCES

1. Yokota, K., Kagawa, S., Shimizu, Y., Akioka, H., Tsurumi, C., Noda, C., Fujimuro, M., Yokosawa, H., Fujiwara, T., Takahashi, E., Ohba, M., Yamasaki, M., DeMartino, G.N., Slaughter, C.A., Toh-e, A. and Tanaka, K. 1996. CDNA cloning of p112, the largest regulatory subunit of the human 26S proteasome, and functional analysis of its yeast homologue, sen3p. *Mol. Biol. Cell* 7: 853-870.
2. Tanaka, K. and Tsurumi, C. 1997. The 26S proteasome: subunits and functions. *Mol. Biol. Rep.* 24: 3-11.
3. Shibahara, T., Kawasaki, H. and Hirano, H. 2004. Mass spectrometric analysis of expression of ATPase subunits encoded by duplicated genes in the 19S regulatory particle of rice 26S proteasome. *Arch. Biochem. Biophys* 421: 34-41.
4. Thompson, H.G., Harris, J.W. and Brody, J.P. 2004. Post-translationally modified S12, absent in transformed breast epithelial cells, is not associated with the 26S proteasome and is induced by proteasome inhibitor. *Int. J. Cancer* 111: 338-347.
5. Tan, Y., Peng, X., Wang, F., You, Z., Dong, Y. and Wang, S. 2006. Effects of tumor necrosis factor- α on the 26S proteasome and 19S regulator in skeletal muscle of severely scalded mice. *J. Burn Care Res.* 27: 226-233.
6. Wang, Z., Aris, V.M., Ogburn, K.D., Soteropoulos, P. and Figueiredo-Pereira, M.E. 2006. Prostaglandin J2 alters pro-survival and pro-death gene expression patterns and 26 S proteasome assembly in human neuroblastoma cells. *J. Biol. Chem.* 281: 21377-21386.
7. Deng, S., Zhou, H., Xiong, R., Lu, Y., Yan, D., Xing, T., Dong, L., Tang, E. and Yang, H. 2007. Over-expression of genes and proteins of ubiquitin specific peptidases (USPs) and proteasome subunits (PSs) in breast cancer tissue observed by the methods of RFDD-PCR and proteomics. *Breast Cancer Res. Treat.* 104: 21-30.
8. Bellizzi, D., Dato, S., Cavalcante, P., Covello, G., Di Cianni, F., Passarino, G., Rose, G. and De Benedictis, G. 2007. Characterization of a bidirectional promoter shared between two human genes related to aging: SIRT3 and PSMD13. *Genomics* 89: 143-150.

CHROMOSOMAL LOCATION

Genetic locus: PSMD13 (human) mapping to 11p15.5.

PRODUCT

PSMD13 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PSMD13 shRNA Plasmid (h): sc-96416-SH and PSMD13 shRNA (h) Lentiviral Particles: sc-96416-V as alternate gene silencing products.

For independent verification of PSMD13 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-96416A, sc-96416B and sc-96416C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PSMD13 siRNA (h) is recommended for the inhibition of PSMD13 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PSMD13 gene expression knockdown using RT-PCR Primer: PSMD13 (h)-PR: sc-96416-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.