



ELP4 siRNA (h): sc-96714

BACKGROUND

In *Saccharomyces cerevisiae*, the hyperphosphorylated form of RNA polymerase II (RNAP II) mediates transcription elongation, and associates with the Elongator complex, which contains six subunits. The Elongator complex can be separated into two subcomplexes; one consisting of Elp1, Elp2 and Elp3, and the other consisting of Elp4, Elp5 and Elp6. The Elongator complex acetylates both core histones and nucleosomal substrates, and directs its activity specifically towards the N-terminal tails of Histone H3 and Histone H4. An analogous complex exists in mammals and contains a variety of proteins that are functional homologs of their yeast counterparts. ELP4 (elongation protein 4), also known as PAX6NEB, is a 424 amino acid protein that localizes to both the cytoplasm and the nucleus and exists as a component of the Elongator complex. Widely expressed as multiple alternatively spliced isoforms, ELP4 is involved in transcriptional regulation and may play a role in chromatin remodeling.

REFERENCES

1. Winkler, G.S., et al. 2001. RNA polymerase II elongator holoenzyme is composed of two discrete subcomplexes. *J. Biol. Chem.* 276: 32743-32749.
2. Hawkes, N.A., et al. 2002. Purification and characterization of the human elongator complex. *J. Biol. Chem.* 277: 3047-3052.
3. Kleinjan, D.A., et al. 2002. Characterization of a novel gene adjacent to PAX6, revealing synteny conservation with functional significance. *Mamm. Genome* 13: 102-107.
4. Kim, J.H., et al. 2002. Human Elongator facilitates RNA polymerase II transcription through chromatin. *Proc. Natl. Acad. Sci. USA* 99: 1241-1246.
5. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 606985. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Li, F., et al. 2004. The Elp4 subunit of human Elongator complex partially complements the growth defects of yeast ELP4 deletion strain. *Yi Chuan Xue Bao* 31: 668-674.

CHROMOSOMAL LOCATION

Genetic locus: ELP4 (human) mapping to 11p13.

PRODUCT

ELP4 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ELP4 shRNA Plasmid (h): sc-96714-SH and ELP4 shRNA (h) Lentiviral Particles: sc-96714-V as alternate gene silencing products.

For independent verification of ELP4 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-96714A, sc-96714B and sc-96714C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ELP4 siRNA (h) is recommended for the inhibition of ELP4 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ELP4 gene expression knockdown using RT-PCR Primer: ELP4 (h)-PR: sc-96714-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.