

TRIM44 siRNA (h): sc-96836

BACKGROUND

The tripartite motif (TRIM) family of proteins are characterized by a conserved TRIM domain that includes a coiled-coil region, a B box-type zinc finger, one RING finger and three zinc-binding domains. TRIM proteins are involved in a wide variety of cellular processes such as cell development, proliferation, differentiation, oncogenesis and apoptosis. Many TRIM proteins are induced by type I and type II interferons, making them crucial for development of pathogen-resistance. TRIM44 (tripartite motif-containing protein 44), also known as Protein DIPB, is a 344 amino acid protein that contains a specific B box-type zinc finger that is found in ubiquitin hydrolases and ubiquitin-specific proteases. TRIM44 inhibits ubiquitination of TRIM17 and may also be involved in the process of neuronal cell differentiation and maturation.

REFERENCES

1. Boutou, E., Matsas, R. and Mamalaki, A. 2001. Isolation of a mouse brain cDNA expressed in developing neuroblasts and mature neurons. *Brain Res. Mol. Brain Res.* 86: 153-167.
2. Leong, P.W., Liew, K., Lim, W. and Chow, V.T. 2002. Differential display RT-PCR analysis of enterovirus-71-infected rhabdomyosarcoma cells reveals mRNA expression responses of multiple human genes with known and novel functions. *Virology* 295: 147-159.
3. Nisole, S., Stoye, J.P. and Saïb, A. 2005. TRIM family proteins: retroviral restriction and antiviral defence. *Nat. Rev. Microbiol.* 3: 799-808.
4. Järvinen, A.K., Autio, R., Kilpinen, S., Saarela, M., Leivo, I., Grenman, R., Mäkitie, A.A. and Monni, O. 2008. High-resolution copy number and gene expression microarray analyses of head and neck squamous cell carcinoma cell lines of tongue and larynx. *Genes Chromosomes Cancer* 47: 500-509.
5. Ozato, K., Shin, D.M., Chang, T.H. and Morse, H.C. 2008. TRIM family proteins and their emerging roles in innate immunity. *Nat. Rev. Immunol.* 8: 849-860.
6. Urano, T., Usui, T., Takeda, S., Ikeda, K., Okada, A., Ishida, Y., Iwayanagi, T., Otomo, J., Ouchi, Y. and Inoue, S. 2009. TRIM44 interacts with and stabilizes terf, a TRIM ubiquitin E3 ligase. *Biochem. Biophys. Res. Commun.* 383: 263-268.
7. McNab, F.W., Rajsbaum, R., Stoye, J.P. and O'Garra, A. 2010. Tripartite-motif proteins and innate immune regulation. *Curr. Opin. Immunol.* 23: 46-56.
8. Peters, C.J., Rees, J.R., Hardwick, R.H., Hardwick, J.S., Vowler, S.L., Ong, C.A., Zhang, C., Save, V., O'Donovan, M., Rassl, D., Alderson, D., Caldas, C. and Fitzgerald, R.C. 2010. A 4-gene signature predicts survival of patients with resected adenocarcinoma of the esophagus, junction, and gastric cardia. *Gastroenterology* 139: 1995-2004.

CHROMOSOMAL LOCATION

Genetic locus: TRIM44 (human) mapping to 11p13.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

TRIM44 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TRIM44 shRNA Plasmid (h): sc-96836-SH and TRIM44 shRNA (h) Lentiviral Particles: sc-96836-V as alternate gene silencing products.

For independent verification of TRIM44 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-96836A, sc-96836B and sc-96836C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TRIM44 siRNA (h) is recommended for the inhibition of TRIM44 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TRIM44 gene expression knockdown using RT-PCR Primer: TRIM44 (h)-PR: sc-96836-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.