

ATP5L siRNA (h): sc-97057

BACKGROUND

Mitochondrial ATP synthase is composed of two multi-subunit complexes that utilize an inner membrane electrochemical gradient to catalyze the synthesis of ATP during oxidative phosphorylation. The two multi-subunit complexes are designated F₁ and F₀, the former of which comprises the soluble catalytic core and the latter of which comprises the membrane-spanning proton channel of ATP synthase. F₁ consists of five distinct subunits, designated ATP5A, ATP5B, ATP5C1, ATP5D and ATP5E, while F₀ consists of ten subunits, designated ATP5H, ATP5G1, ATP5I, ATP5G2, ATP5J2, ATP5J, ATP5G3, ATP5S, ATP5F1 and ATP5L. ATP5L, also known as ATP5JG, is a 103 amino acid protein that localizes to the mitochondrial membrane and exists as a subunit of the F₁ complex.

REFERENCES

1. Elston, T., et al. 1998. Energy transduction in ATP synthase. *Nature* 391: 510-513.
2. Wang, H., et al. 1998. Energy transduction in the F₁ motor of ATP synthase. *Nature* 396: 279-282.
3. Aggeler, R., et al. 2002. A functionally active human F₁F₀ ATPase can be purified by immunocapture from heart tissue and fibroblast cell lines. Subunit structure and activity studies. *J. Biol. Chem.* 277: 33906-33912.
4. Leyva, J.A., et al. 2003. Understanding ATP synthesis: structure and mechanism of the F₁-ATPase (Review). *Mol. Membr. Biol.* 20: 27-33.
5. Oster, G., et al. 2003. Rotary protein motors. *Trends Cell Biol.* 13: 114-121.
6. Cross, R.L. 2004. Molecular motors: turning the ATP motor. *Nature* 427: 407-408.

CHROMOSOMAL LOCATION

Genetic locus: ATP5L (human) mapping to 11q23.3.

PRODUCT

ATP5L siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ATP5L shRNA Plasmid (h): sc-97057-SH and ATP5L shRNA (h) Lentiviral Particles: sc-97057-V as alternate gene silencing products.

For independent verification of ATP5L (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-97057A, sc-97057B and sc-97057C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ATP5L siRNA (h) is recommended for the inhibition of ATP5L expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ATP5L gene expression knockdown using RT-PCR Primer: ATP5L (h)-PR: sc-97057-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.