

PSG4 siRNA (h): sc-97632

BACKGROUND

PSG4 (pregnancy specific β -1-glycoprotein 4), also known as PSG9 or CGM4, is a 419 amino acid secreted protein that exists as 2 alternatively spliced isoforms. PSG4 belongs to the PSG family, a group of closely related secreted glycoproteins that are highly expressed in fetal placental syncytiotrophoblast cells. Members of the PSG protein family have a characteristic N-terminal domain that is homologous to the immunoglobulin variable region. PSGs become detectable in serum during the first two to three weeks of pregnancy and increase as the pregnancy progresses, eventually representing the most abundant fetal protein in the maternal blood at term. PSGs function to stimulate secretion of TH2-type cytokines from monocytes, and they may also modulate the maternal immune system during pregnancy, thereby protecting the semi-allopathic fetus from rejection.

REFERENCES

1. Chan, W.Y., et al. 1988. Characterization of cDNA encoding human pregnancy-specific β 1-glycoprotein from placenta and extraplacental tissues and their comparison with carcinoembryonic antigen. *DNA* 7: 545-555.
2. Zimmermann, W., et al. 1989. cDNA cloning demonstrates the expression of pregnancy-specific glycoprotein genes, a subgroup of the carcinoembryonic antigen gene family, in fetal liver. *Biochem. Biophys. Res. Commun.* 163: 1197-1209.
3. Thompson, J., et al. 1990. The human pregnancy-specific glycoprotein genes are tightly linked on the long arm of chromosome 19 and are coordinately expressed. *Biochem. Biophys. Res. Commun.* 167: 848-859.
4. Barnett, T.R., et al. 1990. Characterization of two new members of the pregnancy-specific β 1-glycoprotein family from the myeloid cell line KG-1 and suggestion of two distinct classes of transcription unit. *Biochemistry* 29: 10213-10218.
5. Chan, W.Y., et al. 1991. Characterization of new members of the pregnancy-specific β 1-glycoprotein family. *Mol. Cell. Biochem.* 106: 161-170.
6. Tynan, K., et al. 1992. Assembly and analysis of cosmid contigs in the CEA-gene family region of human chromosome 19. *Nucleic Acids Res.* 20: 1629-1636.
7. Teglund, S., et al. 1994. The pregnancy-specific glycoprotein (PSG) gene cluster on human chromosome 19: fine structure of the 11 PSG genes and identification of 6 new genes forming a third subgroup within the carcinoembryonic antigen (CEA) family. *Genomics* 23: 669-684.
8. Beauchemin, N., et al. 1999. Redefined nomenclature for members of the carcinoembryonic antigen family. *Exp. Cell Res.* 252: 243-249.
9. Online Mendelian Inheritance in Man, OMIM[™]. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 176393. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: PSG4 (human) mapping to 19q13.31.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

PSG4 siRNA (h) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PSG4 shRNA Plasmid (h): sc-97632-SH and PSG4 shRNA (h) Lentiviral Particles: sc-97632-V as alternate gene silencing products.

For independent verification of PSG4 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-97632A and sc-97632B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PSG4 siRNA (h) is recommended for the inhibition of PSG4 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PSG4 gene expression knockdown using RT-PCR Primer: PSG4 (h)-PR: sc-97632-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.