

ZNF180 siRNA (h): sc-97654

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF180 (zinc finger protein 180), also known as HHZ168, is a 692 amino acid nuclear protein implicated in transcriptional regulation. A member of the Krüppel C₂H₂-type zinc finger protein family, ZNF180 contains one KRAB domain and 12 C₂H₂-type zinc fingers, and undergoes post-translational phosphorylation following DNA damage. The gene encoding ZNF180 maps to human chromosome 19q13.31.

REFERENCES

1. Bray, P., Lichter, P., Thiesen, H.J., Ward, D.C. and Dawid, I.B. 1991. Characterization and mapping of human genes encoding zinc finger proteins. *Proc. Natl. Acad. Sci. USA* 88: 9563-9567.
2. Lichter, P., Bray, P., Ried, T., Dawid, I.B. and Ward, D.C. 1992. Clustering of C₂H₂ zinc finger motif sequences within telomeric and fragile site regions of human chromosomes. *Genomics* 13: 999-1007.
3. Wang, R., Cukerman, E., Heng, H.H. and Liew, C.C. 1996. Identification of a locus of zinc finger genes in human chromosome 19q13.1-q13.3 region by fluorescence *in situ* hybridization. *Somat. Cell Mol. Genet.* 22: 245-248.
4. Shannon, M., Hamilton, A.T., Gordon, L., Branscomb, E. and Stubbs, L. 2003. Differential expansion of zinc-finger transcription factor loci in homologous human and mouse gene clusters. *Genome Res.* 13: 1097-1110.
5. Huntley, S., Baggott, D.M., Hamilton, A.T., Tran-Gyamfi, M., Yang, S., Kim, J., Gordon, L., Branscomb, E. and Stubbs, L. 2006. A comprehensive catalog of human KRAB-associated zinc finger genes: insights into the evolutionary history of a large family of transcriptional repressors. *Genome Res.* 16: 669-677.
6. Filion, G.J., Zhenilo, S., Salozhin, S., Yamada, D., Prokhortchouk, E. and Defossez, P.A. 2006. A family of human zinc finger proteins that bind methylated DNA and repress transcription. *Mol. Cell. Biol.* 26: 169-181.
7. Tian, C.Y., Zhang, L.Q. and He, F.C. 2006. Progress in the study of KRAB zinc finger protein. *Yi Chuan* 28: 1451-1456.
8. Matsuoka, S., Ballif, B.A., Smogorzewska, A., McDonald, E.R., Hurov, K.E., Luo, J., Bakalarski, C.E., Zhao, Z., Solimini, N., Lerenthal, Y., Shiloh, Y., Gygi, S.P. and Elledge, S.J. 2007. ATM and ATR substrate analysis reveals extensive protein networks responsive to DNA damage. *Science* 316: 1160-1166.

CHROMOSOMAL LOCATION

Genetic locus: ZNF180 (human) mapping to 19q13.31.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

ZNF180 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZNF180 shRNA Plasmid (h): sc-97654-SH and ZNF180 shRNA (h) Lentiviral Particles: sc-97654-V as alternate gene silencing products.

For independent verification of ZNF180 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-97654A, sc-97654B and sc-97654C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ZNF180 siRNA (h) is recommended for the inhibition of ZNF180 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZNF180 gene expression knockdown using RT-PCR Primer: ZNF180 (h)-PR: sc-97654-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.