



# Ribosomal Protein S28 siRNA (h): sc-97811

## BACKGROUND

Ribosomes, the organelles that catalyze protein synthesis, are composed of a small subunit (40S) and a large subunit (60S) that consist of over 80 distinct ribosomal proteins. Mammalian ribosomal proteins are encoded by multigene families that contain processed pseudogenes and one functional intron-containing gene within their coding regions. Ribosomal Protein S28 (RPS28) is a 69 amino acid protein which belongs to the ribosomal protein S19P family and localizes to cytoplasm. The gene encoding Ribosomal Protein S28 encodes a ribosomal protein that is part of the 40S subunit and maps to human chromosome 19p13.2. The Ribosomal Protein S28 gene has been found in tumors such as insulinomas and esophageal and colon cancers. Like most ribosomal proteins, Ribosomal Protein S28 exists as multiple processed pseudogenes that are scattered throughout the genome.

## REFERENCES

1. Kitaoka, Y., Olvera, J. and Wool, I.G. 1994. The primary structure of rat Ribosomal Protein S23. *Biochem. Biophys. Res. Commun.* 202: 314-320.
2. Vladimirov, S.N., Ivanov, A.V., Karpova, G.G., Musolyamov, A.K., Egorov, T.A., Thiede, B., Wittmann-Liebold, B. and Otto, A. 1996. Characterization of the human small-ribosomal-subunit proteins by N-terminal and internal sequencing, and mass spectrometry. *Eur. J. Biochem.* 239: 144-149.
3. Louie, D.F., Resing, K.A., Lewis, T.S. and Ahn, N.G. 1996. Mass spectrometric analysis of 40 S ribosomal proteins from rat-1 fibroblasts. *J. Biol. Chem.* 271: 28189-28198.
4. Kenmochi, N., Kawaguchi, T., Rozen, S., Davis, E., Goodman, N., Hudson, T.J., Tanaka, T. and Page, D.C. 1998. A map of 75 human ribosomal protein genes. *Genome Res.* 8: 509-523.
5. Yoshihama, M., Uechi, T., Asakawa, S., Kawasaki, K., Kato, S., Higa, S., Maeda, N., Minoshima, S., Tanaka, T., Shimizu, N. and Kenmochi, N. 2002. The human ribosomal protein genes: sequencing and comparative analysis of 73 genes. *Genome Res.* 12: 379-390.
6. Gauci, S., Helbig, A.O., Slijper, M., Krijgsveld, J., Heck, A.J. and Mohammed, S. 2009. Lys-N and trypsin cover complementary parts of the phosphoproteome in a refined SCX-based approach. *Anal. Chem.* 81: 4493-4501.
7. Malygin, A.A. and Karpova, G.G. 2010. Site-specific cleavage of the 40S ribosomal subunit reveals eukaryote-specific Ribosomal Protein S28 in the subunit head. *FEBS Lett.* 584: 4396-4400.
8. Burkard, T.R., Planyavsky, M., Kaupe, I., Breitwieser, F.P., Bürckstümmer, T., Bennett, K.L., Superti-Furga, G. and Colinge, J. 2011. Initial characterization of the human central proteome. *BMC Syst. Biol.* 5: 17.

## CHROMOSOMAL LOCATION

Genetic locus: RPS28 (human) mapping to 19p13.2.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## PRODUCT

Ribosomal Protein S28 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Ribosomal Protein S28 shRNA Plasmid (h): sc-97811-SH and Ribosomal Protein S28 shRNA (h) Lentiviral Particles: sc-97811-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

Ribosomal Protein S28 siRNA (h) is recommended for the inhibition of Ribosomal Protein S28 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.