SANTA CRUZ BIOTECHNOLOGY, INC.

FXYD7 siRNA (h): sc-97832



BACKGROUND

The mammalian FXYD family maintains Na⁺ and K⁺ gradients between the intracellular and extracellular milieus of cells in processes such as renal Na+reabsorption, muscle contraction and neuronal excitability. FXYDs are singlespan membrane proteins that share a 35 amino acid signature domain, beginning with the sequence PFXYD and containing seven invariant and six conserved amino acids. Members of the FXYD family include phospholemman (FXYD1), FXYD2 (the γ subunit of the Na+/K+-ATPase), FXYD3 (Mat8 or mammary tumor protein), FXYD4 (CHIF) and Dysadherin (FXYD5). FXYD7 (FXYD domain containing ion transport regulator 7) is an 80 amino acid single-pass membrane protein that likely functions as an ion channel. FXYD7 may also have a regulatory function in brain, which is where it is primarily expressed. FXYD7 contains two conserved glycine residues and a transmembrane (TM) domain important for interaction with Na+/K+-ATPase.

REFERENCES

- 1. Crambert, G., Beguin, P., Uldry, M., Monnet-Tschudi, F., Horisberger, J.D., Garty, H. and Geering, K. 2003. FXYD7, the first brain- and isoform-specific regulator of Na,K-ATPase: biosynthesis and function of its posttranslational modifications. Ann. N.Y. Acad. Sci. 986: 444-448.
- 2. Crambert, G., Li, C., Swee, L.K. and Geering, K. 2004. FXYD7, mapping of functional sites involved in endoplasmic reticulum export, association with and regulation of Na,K-ATPase. J. Biol. Chem. 279: 30888-30895.
- 3. Li, C., Grosdidier, A., Crambert, G., Horisberger, J.D., Michielin, O. and Geering, K. 2004. Structural and functional interaction sites between Na,K-ATPase and FXYD proteins. J. Biol. Chem. 279: 38895-38902.
- 4. Geering, K. 2005. Function of FXYD proteins, regulators of Na, K-ATPase. J. Bioenerg. Biomembr. 37: 387-392.
- 5. Li, C., Crambert, G., Thuillard, D., Roy, S., Schaer, D. and Geering, K. 2005. Role of the transmembrane domain of FXYD7 in structural and functional interactions with Na,K-ATPase. J. Biol. Chem. 280: 42738-42743.
- 6. Delprat, B., Bibert, S. and Geering, K. 2006. FXYD proteins: novel regulators of Na,K-ATPase. Med. Sci. 22: 633-638.
- 7. Tipsmark, C.K. 2008. Identification of FXYD protein genes in a teleost: tissue-specific expression and response to salinity change. Am. J. Physiol. Regul. Integr. Comp. Physiol. 294: R1367-R1378.
- 8. Peng, L., Huang, R., Zhang, S. and Hertz, L. 2010. Ouabain binding kinetics and FXYD7 expression in astrocytes and neurons in primary cultures: implications for cellular contributions to extracellular K⁺ homeostasis? Neuron Glia Biol. 6: 127-135.

CHROMOSOMAL LOCATION

Genetic locus: FXYD7 (human) mapping to 19q13.12.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

FXYD7 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see FXYD7 shRNA Plasmid (h): sc-97832-SH and FXYD7 shRNA (h) Lentiviral Particles: sc-97832-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNAse-free water provided. Resuspension of the siRNA duplex in 330 µl of RNAse-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

FXYD7 siRNA (h) is recommended for the inhibition of FXYD7 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor FXYD7 gene expression knockdown using RT-PCR Primer: FXYD7 (h)-PR: sc-97832-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.