

SPRED3 siRNA (h): sc-97920

BACKGROUND

SPRED3 (sprouty-related, EVH1 domain-containing protein 3) is a 410 amino acid peripheral membrane protein belonging to the sprouty/SPRED family of proteins. SPRED3 functions as a tyrosine kinase that regulates the activity of the ERK (also known as MAP kinase) cascade by inhibiting the growth-factor-mediated activation of ERK. SPRED3 is subject to tyrosine phosphorylation and contains one KBD domain, one SPR (sprouty) domain and one WH1 domain. The gene encoding SPRED3 maps to human chromosome 19q13.2, which consists of over 63 million bases, houses approximately 1,400 genes and is recognized for having the greatest gene density of the human chromosomes. It is the genetic home for a number of immunoglobulin (Ig) superfamily members, including the killer cell and leukocyte Ig-like receptors, a number of ICAMs, the CEACAM and PSG families and Fc receptors (FcRs).

REFERENCES

- Mizoguchi, M., Nutt, C.L. and Louis, D.N. 2004. Mutation analysis of CBL-C and SPRED3 on 19q in human glioblastoma. *Neurogenetics* 5: 81-82.
- Lock, P., I, S.T., Straffon, A.F., Schieb, H., Hovens, C.M. and Stylli, S.S. 2006. Spred-2 steady-state levels are regulated by phosphorylation and Cbl-mediated ubiquitination. *Biochem. Biophys. Res. Commun.* 351: 1018-1023.
- Katoh, Y. and Katoh, M. 2006. FGF signaling inhibitor, SPRY4, is evolutionarily conserved target of WNT signaling pathway in progenitor cells. *Int. J. Mol. Med.* 17: 529-532.
- King, J.A., Corcoran, N.M., D'Abaco, G.M., Straffon, A.F., Smith, C.T., Poon, C.L., Buchert, M., I, S., Hall, N.E., Lock, P. and Hovens, C.M. 2006. Evc-3: a liver enriched suppressor of Ras/MAPK signaling. *J. Hepatol.* 44: 758-767.
- Bundschu, K., Walter, U. and Schuh, K. 2007. Getting a first clue about SPRED functions. *Bioessays* 29: 897-907.
- Guy, G.R., Jackson, R.A., Yusoff, P. and Chow, S.Y. 2009. Sprouty proteins: modified modulators, matchmakers or missing links? *J. Endocrinol.* 203: 191-202.
- Spurlock, G., Bennett, E., Chuzhanova, N., Thomas, N., Jim, H.P., Side, L., Davies, S., Haan, E., Kerr, B., Huson, S.M. and Upadhyaya, M. 2009. SPRED1 mutations (Legius syndrome): another clinically useful genotype for dissecting the neurofibromatosis type 1 phenotype. *J. Med. Genet.* 46: 431-437.

CHROMOSOMAL LOCATION

Genetic locus: SPRED3 (human) mapping to 19q13.2.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

SPRED3 siRNA (h) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SPRED3 shRNA Plasmid (h): sc-97920-SH and SPRED3 shRNA (h) Lentiviral Particles: sc-97920-V as alternate gene silencing products.

For independent verification of SPRED3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-97920A and sc-97920B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SPRED3 siRNA (h) is recommended for the inhibition of SPRED3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SPRED3 gene expression knockdown using RT-PCR Primer: SPRED3 (h)-PR: sc-97920-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.