

# 4E-BP1 (P-1): sc-9977

## BACKGROUND

The translation of proteins from eukaryotic mRNA is initiated by the multisubunit complex eIF-4F, which associates with the mRNA 5' cap structure. eIF-4E, a component of eIF-4F, is responsible for binding to the 5' cap structure and for the assembly of the eIF-4F complex. The regulatory protein 4E-BP1, also referred to as PHAS-I, inhibits eIF-4E function. Phosphorylation of 4E-BP1 by S6 kinase p70, MAP kinases or PKCs causes the disassociation of 4E-BP1 from eIF-4E, promoting translation. A protein that is functionally related to 4E-BP1, designated 4E-BP2, also associates with eIF-4E.

## CHROMOSOMAL LOCATION

Genetic locus: EIF4EBP1 (human) mapping to 8p11.23; Eif4ebp1 (mouse) mapping to 8 A2.

## SOURCE

4E-BP1 (P-1) is a mouse monoclonal antibody raised against full length 4E-BP1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

4E-BP1 (P-1) is available conjugated to agarose (sc-9977 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-9977 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-9977 PE), fluorescein (sc-9977 FITC), Alexa Fluor® 488 (sc-9977 AF488), Alexa Fluor® 546 (sc-9977 AF546), Alexa Fluor® 594 (sc-9977 AF594) or Alexa Fluor® 647 (sc-9977 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-9977 AF680) or Alexa Fluor® 790 (sc-9977 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

## APPLICATIONS

4E-BP1 (P-1) is recommended for detection of 4E-BP1 of mouse, rat, human and porcine origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for 4E-BP1 siRNA (h): sc-29594, 4E-BP1 siRNA (m): sc-29595, 4E-BP1 shRNA Plasmid (h): sc-29594-SH, 4E-BP1 shRNA Plasmid (m): sc-29595-SH, 4E-BP1 shRNA (h) Lentiviral Particles: sc-29594-V and 4E-BP1 shRNA (m) Lentiviral Particles: sc-29595-V.

Molecular Weight of 4E-BP1: 21 kDa.

Positive Controls: A-431 whole cell lysate: sc-2201, K-562 whole cell lysate: sc-2203 or KNRK whole cell lysate: sc-2214.


## RESEARCH USE

For research use only, not for use in diagnostic procedures.


## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## DATA



4E-BP1 (P-1): sc-9977. Western blot analysis of 4E-BP1 expression in A-431 (A), K-562 (B), NIH/3T3 (C) and KNRK (D) whole cell lysates.



4E-BP1 (P-1): sc-9977. Immunoperoxidase staining of formalin fixed, paraffin-embedded human pancreas tissue showing cytoplasmic staining of glandular cells.

## SELECT PRODUCT CITATIONS

- Xu, G., et al. 2004. Pharmacogenomic profiling of the PI3K/PTEN-Akt-mTOR pathway in common human tumors. *Int. J. Oncol.* 24: 893-900.
- Zheng, W., et al. 2016. Far upstream element-binding protein 1 binds the 3' untranslated region of PKD2 and suppresses its translation. *J. Am. Soc. Nephrol.* 27: 2645-2657.
- Lu, J., et al. 2017. Tunneling nanotubes promote intercellular mitochondria transfer followed by increased invasiveness in bladder cancer cells. *Oncotarget* 8: 15539-15552.
- Wang, X., et al. 2017. Sestrin2 and sestrin3 suppress NK-92 cell-mediated cytotoxic activity on ovarian cancer cells through AMPK and mTORC1 signaling. *Oncotarget* 8: 90132-90143.
- Nie, X., et al. 2018. mTOR acts as a pivotal signaling hub for neural crest cells during craniofacial development. *PLoS Genet.* 14: e1007491.
- Sanguesa, G., et al. 2018. Chronic fructose intake does not induce liver steatosis and inflammation in female Sprague-Dawley rats, but causes hypertriglyceridemia related to decreased VLDL receptor expression. *Eur. J. Nutr.* 58: 1283-1297.
- Lee, M.K., et al. 2018. *Pyropia yezoensis* protein supplementation prevents dexamethasone-induced muscle atrophy in C57BL/6 mice. *Mar. Drugs* 16 pii: E328.
- Li, X., et al. 2018. Hypoxia-induced autophagy of stellate cells inhibits expression and secretion of lumican into microenvironment of pancreatic ductal adenocarcinoma. *Cell Death Differ.* 26: 382-393.
- Liu, P., et al. 2019. Wound healing potential of spirulina protein on CCD-986sk cells. *Mar. Drugs* 17 pii: E130.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.